Feb. 26. 2018

Spintronics Basic II

Hisazumi Akai

The Institute for Solid State Phisucs

The University of Tokyo

Contents

- Origin of magnetism and magnetic structures
  - Exchange interaction
  - Exchange interaction between two atoms
  - Heitler-London limit and molecular orbital limit
  - Direct exchange and kinetic exchange
  - Atoms in a crystal
  - Kanamori-Goodenough rule
  - Double exchange
  - Magnetic anisotropy
  - Anisotropic exchange
  - Dzialshinsky-Moriya interaction
  - Heisenberg model

Interaction between electrons

■ two electrons –Pauli's exclusion principle



possibility that two electrons come closer
→ increase in coulomb energy



less probability of two electrons meet
→ suppress increase in coulomb energy

parallel spin is preferable

Interaction between electrons

■ two electrons –Pauli's exclusion principle



possibility that two electrons come closer
→ increase in coulomb energy



less probability that two electro meet
→ suppress increase in coulomb energy

reason why electrons do not come closer is not coulomb repulsion but Pauli's exclusion principle

parallel spin is preferable

Kinetic energy of electrons

two electrons –Pauli's exclusion principle



no limitation to the volume where electrons can move freely
→ no increase in the kinetic energy



volume where electrons can move is halved
→ increase in zero-point energy

$$\Delta r \Delta p \sim \hbar$$

antiparallel spin is preferable

Exchange interaction between electrons

- energy difference between parallel and antiparallel couples of electrons

E_X = E  − E 

Single atom

- electrons are bounded in a potential
- neutral atom with an odd number of electrons
  - nonzero total electron spin
  - nonzero angular momentum
  - shows magnetism (paramagnetic)



Tow atoms

- interaction between atomic magnetic moments
  - parallel or antiparallel coupling
  - exchange interaction between atoms

$E_X = E$  $- E$ 

parallel if $E_X > 0$, antiparallel otherwise

Parallel or antiparallel?

- Simple model I Heitler–London limit
  - two hydrogen atoms (hydrogen atom1 and hydrogen atom2)
  - potential of each atom $v(r-R)$

	atomic hamiltonian	atomic orbital
hydrogen atom1	$H_1 = -\nabla_1^2 + v(r_1 - R_1)$	$H_1\phi_1 = \epsilon\phi_1$
hydrogen atom2	$H_2 = -\nabla_2^2 + v(r_2 - R_2)$	$H_2\phi_2 = \epsilon\phi_2$

（we use Slater's atomic unit: $\hbar = 1,\ m = 1/2,\ e^2 = 2$ ）

Parallel or antiparallel?

Simple model I: Heitler–London limit

- assume that each atom keeps charge neutrality
- assume that the wave function of hydrogen molecule is a product of 1s wave functions ϕ's of two hydrogen atoms

hamiltonian of hydrogen molecule

coulombic interaction

$$H = H_1 + H_2 + H_{12} + V(r_1 - r_2)$$

$$H_{12} = v(r_1 - R_2) + v(r_2 - R_1)$$

atractive potential due to the other atom

molecular wavefunction $\Psi(r_1, r_2) \sim \phi_1(r_1)\phi_2(r_2)$

molecular energy $E \simeq \int \Psi^* H \Psi \, d^3r$

Parallel or antiparallel?

- Simple model I: Heitler–London limit
  - antisymmetrization (sign change associated with exchange of two electrons)
  - two cases depending on spin configuration

s=0 state (spin singlet)



$$\Psi(r_1, r_2) = A_0\{\phi_1(r_1)\phi_2(r_2) + \phi_1(r_2)\phi_2(r_1)\}$$

(symmetric orbital, antisymmetric spin)

s=1 state (spin triplet)



$$\Psi(r_1, r_2) = A_1\{\phi_1(r_1)\phi_2(r_2) - \phi_1(r_2)\phi_2(r_1)\}$$

(antisymmetric orbital, symmetric spin)

Parallel or antiparallel?

Simple model I: Heitler–London limit

expectation value of energy

spin singlet

$$\int \Psi_0(r_1, r_2)(H_1 + H_2 + H_{12} + V)\Psi_0(r_1, r_2)dr_1dr_2 = E_0$$

spin triplet

$$\int \Psi_1(r_1, r_2)(H_1 + H_2 + H_{12} + V)\Psi_1(r_1, r_2)dr_1dr_2 = E_1$$

$$E_{\mathrm{x}} = E_0 - E_1$$

Parallel or antiparallel?

Simple model I: Heitler–London limit

s=0 state (spin singlet)

$$\Psi_0(r_1, r_2) = A_0\{\phi_1(r_1)\phi_2(r_2) + \phi_1(r_2)\phi_2(r_1)\}$$

$$\Psi_0(r_1, r_2) \neq 0 \qquad \text{even if} \quad r_1 = r_2$$

large interaction V



s=1 state (spin triplet)

$$\Psi_1(r_1, r_2) = A_1\{\phi_1(r_1)\phi_2(r_2) - \phi_1(r_2)\phi_2(r_1)\}$$

$$\Psi_1(r_1, r_2) = 0 \qquad \text{if} \qquad r_1 = r_2$$

small interaction V

$$\boxed{\int \Psi_0(r_1, r_2) V \Psi_0(r_1, r_2) dr_1 dr_2 > \int \Psi_1(r_1, r_2) V \Psi_1(r_1, r_2) dr_1 dr_2}$$

Parallel or antiparallel?

Simple model I: Heitler–London limit

s=0 state（spin singlet）

$$\Psi_0(r_1, r_2) = A_0\{\phi_1(r_1)\phi_2(r_2) + \phi_1(r_2)\phi_2(r_1)\}$$

$$\int \Psi_0(r_1, r_2)(H_1 + H_2 + H_{12})\Psi_0(r_1, r_2)dr_1dr_2 = 2\epsilon'$$

s=1 state（spin triplet）

$$\Psi_1(r_1, r_2) = A_1\{\phi_1(r_1)\phi_2(r_2) - \phi_1(r_2)\phi_2(r_1)\}$$

$$\int \Psi_1(r_1, r_2)(H_1 + H_2 + H_{12})\Psi_1(r_1, r_2)dr_1dr_2 = 2\epsilon'$$

the same energy except coulomb interaction energy

$$\left(\epsilon' = \epsilon + \int \phi_1(r_1)v(r - R_1)\phi_1(r_1)dr_1 < \epsilon\right)$$

Heitler–London limit

therefore

$$\int \Psi_0(r_1, r_2) H \Psi_0(r_1, r_2) dr_1 dr_2 > \int \Psi_1(r_1, r_2) H \Psi_1(r_1, r_2) dr_1 dr_2$$

$$E_{\mathrm{x}} = E_0 - E_1 > 0$$

- triplet state realized in Heitler-London limit
- not corresponding to most two-atom molecules
  - two-atom molecules are mostly spin singlet
- molecular bonding is caused by H_{12}
  - ε' < ε : effect of attractive potential of neighbor

Parallel or antiparallel?

- Simple model 2: molecular orbital limit
  - two hydrogen atoms (hydrogen atom1 and hydrogen atom2)
  - wavefunctions are molecular orbitals constructed from two atomic orbitals
  - two molecular orbitals: bonding and antibonding states
  - how to place two electrons in these molecular orbitals?

bonding state

$$\Psi_{\mathrm{b}}(r) = A\{\phi_1(r) + \phi_2(r)\}$$

antibonding state

$$\Psi_{\mathrm{a}}(r) = A\{\phi_1(r) - \phi_2(r)\}$$

$$\left(\quad \phi_i(r) \equiv \phi(r - R_i) \quad \right)$$

Parallel or antiparallel?

Simple model 2: molecular orbital limit

antibonding state

$$\Psi_{\mathrm{a}}(r) = A\{\phi_1(r) - \phi_2(r)\}$$

single electron energy



most stable if coulombic interaction is missing

bonding state

$$\Psi_{\mathrm{b}}(r) = A\{\phi_1(r) + \phi_2(r)\}$$

Parallel or antiparallel?

Simple model 2: molecular orbital limit

antibonding state

$$\Psi_{\mathrm{a}}(r) = A\{\phi_1(r) - \phi_2(r)\}$$



bonding state

$$\Psi_{\mathrm{b}}(r) = A\{\phi_1(r) + \phi_2(r)\}$$

Parallel or antiparallel?

Simple model 2: molecular orbital limit



$E_0 < E_1$ or $E_0 > E_1$?

Hitler–London vs. molecular orbital limits

- Heitler–London limit excludes ionic states
  - strong correlation limit
  - electron interaction energy $\gg$ kinetic energy
- equal probability of ionic and neutral states in molecular orbital limit
  - weak correlation limit
  - electron interaction energy $\ll$ kinetic energy



Analysis using simple model

2-site Hubbard model

$$H = t \sum_{\sigma} (a_{1\sigma}^{\dagger} a_{2\sigma} + a_{2\sigma}^{\dagger} a_{1\sigma}) + U \sum_{i=1,2} n_{i\uparrow} n_{i\downarrow}$$



Exact solution

- ground state is spin singlet state for any U



Exact solution

- ground state at U=0 and ∞



Mean field approximation (MFA)

- nonmagnetic to ferromagnetic transition occurs



Mean field approximation (MFA)

- nonmagnetic to ferromagnetic transition occurs



In reality

- somewhere in between Heitler-London and molecular orbital
  - starting from Heitler-London
    - ionic states gradually mixed into neutral states
    - although ionic states have higher energies, energy gain due to electron transfer is expected
  - starting from molecular orbital limit
    - ionic states are suppress owing to their higher energy
    - bonding–antibonding splitting decreases, which reduce energy gain due to electron transfer

real atomic exchange couplings

- between two atomic spins $\boldsymbol{S}_1$ and $\boldsymbol{S}_2$ of atom 1 and atom 2, there exists an interaction of the form

$$E_{\mathrm{x}} = E_0 - E_1 = -2(J_{12} + W_{12})\boldsymbol{S}_1 \cdot \boldsymbol{S}_2$$

where

$$J_{12} > 0, \quad W_{12} < 0$$

is always satisfied and in many cases

$$J_{12} + W_{12} < 0$$

Direct and kinetic exchanges

- J: direct exchange
- W: kinetic exchange
- in many cases, W determines magnetic structures

Hereafter, we use J to indicate a sum of J and W

$$J_{12} + W_{12} \Rightarrow J_{12}$$

Therefore J_{12} can be either positive or negative.
A positive J is ferromagnetic and vice versa.

If many atoms exist

- when interaction is strong enough, spontaneous symmetry breaking occurs
  - magnetic moment arises at each site
  - stable magnetic structure realized
  - in this situation MFA gives reasonable descriptions
  - energy of systems depends of magnetic structures
  - lowest energy structure is realized at ground states

Magnetic ions in solids

- consider a situation where interactions are large enough to realize a magnetic state
  - each atom carries a magnetic moment
- consider at T=0
  - which magnetic structure is most stable?
  - what determines the magnetic structure?

Kanamori–Goodenough rule

- exchange interactions of oxides and haleids

180° configuration



$J_{12} < 0$



$J_{12} > 0$

90° configuration



$J_{12} > 0$



$J_{12} < 0$

Analysis using Hubbard model

- simple but qualitatively correct

$$H = t\sum_{\sigma}(a_{1\sigma}^{\dagger}a_{2\sigma} + a_{2\sigma}^{\dagger}a_{1\sigma}) + U\sum_{i=1,2} n_{i\uparrow}n_{i\downarrow}$$

superexchange

energy gain due to electron transfer $-\dfrac{t^2}{U}$



antiferromagnetic

atom1 atom2

no energy gain because electron transfer is prohibited



ferromagnetic

atom1 atom2

Analysis using Hubbard model

- simple but qualitatively correct

$$H = t\sum_{\sigma}(a_{1\sigma}^{\dagger}a_{2\sigma} + a_{2\sigma}^{\dagger}a_{1\sigma}) + U\sum_{i=1,2} n_{i\uparrow}n_{i\downarrow}$$

superexchange

double-exchange

energy gain due to electron transfer $-\dfrac{t^2}{U}$

energy gain due to electron transfer $-nt$

when holes exist



antiferromagnetic

atom1 atom2



ferromagnetic

atom1 atom2

Ferromagnetism appears if carriers exist

- at half-filled only superexchange works
  - antiferromagnetic (effects in second order of t)
- if carriers exist, double-exchange works in proportion to carrier concentration (effects in first order in t)
  - when n exceeds some value, ferromagnetism appears
- for small t's, even a small n realizes ferromagnetism

$$\frac{t^2}{U} < nt \quad \text{for small} \quad t$$

In terms of band electron

two mechanisms in band picture

ferrimagnetism stabilized

band broadening by $2t$



double-exchange

antiferromagnetism stabilized

widening of split by $2\frac{t^2}{U}$



spuperexchange

Spin-orbit interaction

- a relativistic effect
- in view of electrons turning around a nucleus, the nucleus is turning around the electrons
- turning nucleus thus produces electric current



$$B \propto \frac{ZeL\hbar}{mr^3}$$

$$H_{\mathrm{so}} = \lambda \boldsymbol{L} \cdot \boldsymbol{S}$$

Effects of spin-orbit interaction

- magnetic anisotropy
  - without spin-orbit interaction, spins do not see lattices
  - because of spin-orbit interactions, correlation between directions of spins and lattice orientations arises
  - spins prefer special direction relative to a lattice
- anisotropic exchange interaction
- anti-symmetric exchange interaction (Dzyaloshinsky–Moriya interaction: DMI)
- orbital magnetic moment in crystals

Magnetic anisotropy

- there exists a special direction to which magnetization is apt to aligns





Anisotropic exchange

- orbitals are distorted as a result of spin-orbit interaction
- exchange interaction thus depends on the direction of spins

$$J_{\mathrm{A}} \neq J_{\mathrm{B}}$$

This interaction can be expressed as $\boldsymbol{S}_A \mathbf{J} \boldsymbol{S}_B$ using symmetric tensor $\mathbf{J}$

Dzyaloshinsky–Moriya interaction

- because of spin-orbit interaction, J depends on direction (along or opposite way



Due to sin θdependence, the two spins cant

- from A there is C to the right of interaction line
- from B there is C to the left of interaction line

As a result $J_{\mathrm{AB}} \neq J_{\mathrm{BA}}$

In general J is expressed as

$$J = J_{\mathrm{s}} \pm J_{\mathrm{a}}$$

This can either be expressed using spin as

$$-2J\boldsymbol{S}_A \cdot \boldsymbol{S}_B + \boldsymbol{D} \cdot (\boldsymbol{S}_A \times \boldsymbol{S}_B)$$

The second terms is called anti-symmetric exchange interaction (Dzyalonshinsky–Moriya interaction: DMI). The interaction depends on the angle between direction of two spinsθ as sinθ.

Heisenberg model

- phenomenological model describing magnetism of matter
- describes magnetic behavior of magnetic insulators rather well
- sometimes can be used for phenomenological description of metallic magnetism as well (there also are many cases where this does not work at all)

$$H = -\sum_{<ij>} 2J_{ij}\boldsymbol{S}_i \cdot \boldsymbol{S}_j$$

J_{ij} obtained from electronic structure calculation

- example of calculated exchange coupling constants



Heisenberg model

- magnetic anisotropy, anisotropic exchange, antisymmetric exchange, etc. can be also treated in the framework
- variations such as Ising model, XY model
- typical playground for statistical physics is used to discuss finite temperature properties and phase transitions of magnetic materials.

$$H = -\sum_{<ij>} 2J_{ij}\boldsymbol{S}_i \cdot \boldsymbol{S}_j$$

Summary

- magnetism of insulators in mind, we discussed
  - atomic magnetism
  - molecular magnetism and exchange interactions
  - mechanisms of exchange interactions
  - exchange interactions between magnetic ions in crystals
  - effects of spin-orbit interactions
  - Heisenberg model
- for further study:
  - J. Kanamori, "Magnetism" (Japanese) (Baifukan, Physics Series 7)